arXiv:1212.5450v2 [cond-mat.quant-gas] 8 Feb 2013

Inhomogeneous Superfluid Phases in the Unitary ^{6}Li-^{40}K Mixture.

J.E. Baarsma* and H.T.C. Stoof
Institute for Theoretical Physics, Utrecht University,
Leuvenlaan 4, 3584 CE Utrecht, The Netherlands

We have determined the complete phase diagram of the ^{6}Li-^{40}K mixture within mean-field theory taking the inhomogeneities of the fermion states into account exactly. We find two different inhomogeneous superfluid phases in mixtures with a majority of ^{40}K atoms, namely the Larkin-Ovchinnikov (LO^1) phase with one inhomogeneous direction and a cubic phase (LO^3) where three spatial translational symmetries are broken. We determine the transition between these two phases by solving the Bogoliubov-de Gennes equations in the LO^1 phase. Subsequently, we calculate the atomic density modulation of the atoms in the LO^1 phase and show that for the minority atoms it is sufficiently large to be visible in experiment.

PACS numbers: 03.75.-b, 67.40.-w, 39.25.+k

Since the first realization of superfluidity in a two-component gas of fermionic atoms a large number of exciting experiments have been performed. For instance, with the use of a Feshbach resonance the inter-particle interactions can be tuned and hereby the crossover between the Bose-Einstein condensation of molecules and Cooper pairs was studied [1, 2]. By inducing spin flips in a Fermi gas the number of particles in different spin states can be changed, opening up the possibility of studying the influence of a population imbalance on the phase transition to a superfluid state. This possibility has indeed been materialized in a two-component gas of ^{6}Li atoms and the phase diagram turns out to be governed by a tricritical point below which the gas phase separates into a superfluid and normal region [3–6].

One of the exotic states of matter that may also be realized in a system of ultracold Fermi atoms is a superfluid where the Cooper pairs have a nonzero center-of-mass momentum. The first to propose this possibility in the context of superfluid films in a magnetic field were Fulde and Ferrel [7] and independently Larkin and Ovchinnikov (LO) [8]. Signatures of these so called FFLO phases have been seen in an atomic gas in one spatial dimension [9], but a decisive experiment that observes the FFLO correlations has not been carried out yet. In three spatial dimensions a phase with nonzero momentum Cooper pairs is predicted to be present in the population imbalanced Fermi gas with weak interactions [10–14]. However, the transition temperatures in a weakly-interacting Fermi gas are very low and are expected to be out of reach with present cooling techniques.

In the three-dimensional strongly interacting mixture of ^{6}Li and ^{40}K atoms a Lifshitz instability towards a phase with nonzero momentum Cooper pairs is present at an attainable temperature [15, 16], but the precise form of this superfluid phase has not yet been determined. In this Letter we show that at temperatures below the Lifshitz point the Fermi gas will form the exotic LO phase, which shares the properties of both a solid and a superfluid and is therefore a special kind of supersolid. This is a very exciting prospect, because even though this new kind of superfluid was first proposed by Larkin and Ovchinnikov already in 1964 [8], it has never been observed in a three-dimensional fluid. In the ^{6}Li-^{40}K mixture important steps have already been made experimentally. The Feshbach resonances of the mixture were known for some time [17, 18], and now also expansion under the influence of strong interactions from an optical dipole trap has been realized [19], a large atom number dual-species magneto-optical trap was built [20], and the repulsive polaron has been studied in this mixture [21]. At present, therefore, an ultra-cold ^{6}Li-^{40}K mixture at unitarity is to the best of our knowledge the only system available in the laboratory to observe such a three-dimensional supersolid and study its properties in detail with the accuracy of atomic physics.



FIG. 1: Phase diagram for the unitary mixture of ^{6}Li and ^{40}K atoms (a), where the cross denotes the point for which we calculated the atomic densities in Fig.3. The grey rectangle on the horizontal axis sets the scale for panel (b), where along the line of phase transitions to an inhomogeneous superfluid is shown. Also shown are the Fermi wavevectors of the ^{6}Li(↑) and ^{40}K(↓) atoms and their difference, all at $T = 0$.

The LO state is also expected to be present in a number of other condensed-matter systems, such as neutron stars [22] and heavy-fermion systems [23]. In the context of colorsuperconductivity a large body of theoretical work has been concerned with different inhomogeneous phases [24–28], all considering weak coupling and using Ginzburg-Landau expansions of the free energy. Although continuous (second-order) instabilities can be determined exactly using this expansion, the downfall is that ruling out discontinuous (first-order) transitions is impossible and one is moreover forced to remain close to the normal-superfluid transition.

---

*Electronic address: J.E.Baarsma@uu.nl

In this Letter we calculate for different inhomogeneous superfluids the full inhomogeneous thermodynamic potential within mean-field theory, with which both continuous and discontinuous phase transitions can be described. The Lifshitz point found using this mean-field theory remains present when adding screening and selfenergy effects and its position only changes quantitatively [15, 16]. Adding these effects for the mass-balanced case reproduces the experimentally found phase diagram [16], from which it can be concluded that for these purposes the mean-field calculation already contains the relevant physics, even at unitarity.

We determine the phase diagram of the mixture of ^{6}Li-^{40}K atoms for a majority of ^{40}K atoms, see Fig.1(a), and especially what the structure of the inhomogeneous superfluid is. For a majority of ^{6}Li atoms it was shown already that a tricritical point is present, below which a discontinuous phase transition towards a homogeneous superfluid, and thus phase separation, occurs [15]. We find for small negative polarizations $P = (n_\uparrow - n_\downarrow)/(n_\uparrow - n_\downarrow)$, where $n_{\uparrow(\downarrow)}$ denotes the density of ^{6}Li(^{40}K) atoms, a continuous transition from the normal state (N) to a homogeneous superfluid (HSF), up to the Lifshitz point (LP). For larger majorities of ^{40}K atoms there is a continuous phase transition to an inhomogeneous superfluid phase where translational symmetry is broken in one spatial direction (LO^1) and for even larger population imbalances the continuous transition is to a phase where the translational symmetry is broken in three directions (LO^3). In Fig.1(a) the temperature T is scaled by the critical temperature $T_{c,0}$ at $P = 0$ and the polarization P is scaled by the critical polarization $|P_{c,0}|$ at $T = 0$, because interaction effects in first instance only shift the location of the Lifshitz point [15, 16].

In addition, after explaining how we obtained the above phase diagram, we determine the densities for the two atomic species in the LO^1 phase at the conditions of the cross in Fig.1(a), which are now position dependent with a modulation that can be as big as 10% over one period. This is therefore a convenient signature for the LO^1 phase in experiment, but also other proposals to observe this exotic state of matter have been made [29, 30].

Thermodynamic Potential.— We calculate the grand-canonical thermodynamic potential Ω for the two-component Fermi gas with mass and population imbalance from the microscopic action

$$S = \int d\mathbf{x}d\tau \left[\sum_{\sigma=\uparrow,\downarrow} \phi^*_\sigma \left(\hbar\frac{\partial}{\partial\tau} - \frac{\hbar^2\nabla^2}{2m_\sigma} - \mu_\sigma \right) \phi_\sigma + \frac{|\Delta|^2}{V_0} + \phi^*_\uparrow\phi^*_\downarrow\Delta + \Delta^*\phi_\downarrow\phi_\uparrow \right], \quad (1)$$

where $\Delta(\mathbf{x},\tau)$ is the complex pairing field and $\phi_\sigma(\mathbf{x},\tau)$ are fermionic fields with m_σ the mass and μ_σ the chemical potential of the fermionic atoms. The last two terms in the above action represent respectively the annihilation and creation of a Cooper pair, here consisting of a ^{6}Li and a ^{40}K atom. The microscopic interaction strength V_0 in the above action can be replaced in favor of the effective interaction proportional to the s-wave scattering length a [16], which can be controlled using a Feshbach resonance. In this Letter we always consider unitarity, where $|a| = \infty$. From the action the partition function can be calculated, $Z = \int d[\phi^*]d[\phi]d[\Delta^*]d[\Delta] \exp(-S[\phi,\Delta]/\hbar)$, and subsequently the thermodynamic potential can be obtained via $\Omega = -k_BT \log Z$, where k_BT is the thermal energy with k_B Boltzmann's constant. Densities can in turn be calculated from the thermodynamic potential via $n_\sigma = -\partial\Omega/\partial\mu_\sigma$. In the partition function Z the path integral over the fermionic fields ϕ can be evaluated, after which only the path integral over the pairing field Δ is left with in the integrand an exponent of an effective action $S^{\text{eff}}[\Delta]$. In the mean-field approximation the pairing field is now replaced by its most probable value and the remaining path integral omitted. In practice the thermodynamic potential is then obtained by making an ansatz for the pairing field and thus for the superfluid phase. The thermodynamic potential describing the phase transition from a normal gas to a homogeneous superfluid is calculated by taking for the pairing field $\Delta(\mathbf{x},\tau) = \Delta_0$. Fulde and Ferrel considered a plane wave, $\Delta(\mathbf{x},\tau) = \Delta_0 e^{i\mathbf{q}\cdot\mathbf{x}}$, in which case the Cooper pairs have nonzero momentum $\mathbf{q}$ but the superfluid density is still position independent and thus it is a homogeneous phase. Larkin and Ovchinnikov assumed a standing wave, $\Delta(\mathbf{x},\tau) = \Delta_0 \cos(\mathbf{q}\cdot\mathbf{x})$. In this case not only the Cooper-pair phase but also the superfluid density becomes position dependent, which turns out to be always more favorable than the plane wave ansatz. In the latter case the superfluid is truly inhomogeneous. Obviously, many other choices for the Cooper-pair wavefunction can be made, depending on the situation of interest.

In the case of a position-dependent pairing field the thermodynamic potential not only depends on the amplitude Δ_0 but also on the lattice wavevector $|\mathbf{q}| = q$ of the order parameter, $\Omega(\Delta_0, q)$. For all ansatzes we made for the order parameter we find at the same chemical potential and temperature a continuous phase transition to the superfluid phase. Only after comparison of the different thermodynamic potentials for different pairing fields the most favorable configuration of the superfluid is found. In Fig.2 we show the thermodynamic potentials for different Cooper-pair ansatzes along the line of



FIG. 2: The thermodynamic potential along the normal to superfluid transition line from Fig.1(a). At first the LO^1 (solid line) gives the lowest thermodynamic potential, while for larger polarizations this is true for the LO^3 phase (dashed line). An LO^2 phase (dotted) or more complicated configurations (dashed-dotted and dashed-double-dotted) never form the equilibrium state of this system.

phase transitions to the inhomogeneous superfluid phase. It can be seen that, inside the inhomogeneous superfluid, first the phase with only one broken translational symmetry, the LO^1 ansatz, gives the lowest thermodynamic potential. Further along the line of phase transitions it becomes more favorable to break translational symmetry in three spatial dimensions (LO^3), corresponding to the following ansatz for the order parameter

$$\Delta_3(\mathbf{x}) = \Delta_0\left[\cos(qx) + \cos(qy) + \cos(qz)\right]. \tag{2}$$

Other phases are never favorable. By comparing thermodynamic potentials for different lattice structures we completed the phase diagram for the Fermi mixture of ^{6}Li and ^{40}K atoms, see Fig.1(a). The lattice wavevector, i.e., the value of q for which the thermodynamic potential $\Omega(\Delta_0, q)$ has a global minimum, along the line of the phase transition is shown in Fig.1(b). It can be seen that for $T = 0$ it differs from the difference in Fermi wavevectors, which is due to the fact that we are in the strongly-interacting regime. We found that inside the LO^1 superfluid phase the global minimum of the thermodynamic potential shifts continuously from a nonzero to a zero lattice wavevector q, in which case the superfluid is homogeneous again. This criterion for the transition between the inhomogeneous and homogeneous superfluid is equivalent to the vanishing of the domain-wall energy [11, 14].

LO^1 Green's Function.— We found that for one point on the normal to superfluid transition line the thermodynamic potentials for the LO^1 and the LO^3 phase are equal, see Fig.1(a) and Fig.2. Within the superfluid region the transition line between these two different inhomogeneous phases can be obtained by comparing thermodynamic potentials, but now that we know which phases to compare we can also calculate the transition line more directly. In order to do so, we split the order parameter for the LO^3 phase

$$\Delta(\mathbf{x}) = \Delta_{\mathrm{LO}}(z) + \Delta_\perp[\cos(qx) + \cos(qy)], \tag{3}$$

where $\Delta_{\mathrm{LO}}(z) = \Delta_0\cos(qz)$. We now expand the thermodynamic potential for the cubic phase in the perpendicular order parameter $\Delta_\perp$

$$\Omega_3 \simeq \Omega(\Delta_0, q) + \alpha_\perp(\Delta_0, q)|\Delta_\perp|^2. \tag{4}$$

At the moment $\alpha_\perp$ changes sign the minimum of Ω_3 shifts to a non-zero value of $\Delta_\perp$, which means the LO^3 phase is more favorable than the LO^1 phase. In order to calculate $\alpha_\perp$ we need to determine the Green's function for the LO^1 phase, since

$$\alpha_\perp = -\frac{V}{V_0} + \frac{k_BT}{2|\Delta_\perp|^2}\mathrm{Tr}\left[\mathbf{G}_{\mathrm{LO}}\boldsymbol{\Delta}_\perp\mathbf{G}_{\mathrm{LO}}\boldsymbol{\Delta}_\perp\right], \tag{5}$$

where V is the volume and $\boldsymbol{\Delta}_\perp(x,y) = (\Delta_\perp/\hbar)[\cos(qx) + \cos(qy)]\sigma_x$, with σ_x the first Pauli matrix in Nambu space. The trace in the above equation is taken over real space, imaginary time and Nambu space. The inverse Green's function $\mathbf{G_{LO}}^{-1}$ is known from Eq.(1) and can be expanded in its energy eigenmodes, which are the Bogoliubov quasiparticle wavefunctions. In this manner we obtain the Green's function in the LO^1 phase, which contains the inhomogeneities of this phase in an exact way. The quasiparticle wavefunctions for the Larkin-Ovchinnikov phase we calculate by solving the Bogoliubov-de Gennes equation

$$\begin{pmatrix} -\frac{\hbar^2\nabla^2}{2m_\uparrow} - \mu_\uparrow & \Delta_{\mathrm{LO}}(z) \\ \Delta_{\mathrm{LO}}(z) & \frac{\hbar^2\nabla^2}{2m_\downarrow} + \mu_\downarrow \end{pmatrix}\begin{pmatrix} u_n(\mathbf{x}) \\ v_n(\mathbf{x}) \end{pmatrix} = \hbar\omega_n\begin{pmatrix} u_n(\mathbf{x}) \\ v_n(\mathbf{x}) \end{pmatrix}, \tag{6}$$

where u and v are the quasiparticle coherence factors and n is the band index labeling the energy eigenmode. The inverse Green's function, on the left-hand side of the equation, contains the LO^1 order parameter. Since this pairing field has a periodicity the quasiparticle wavefunctions are according to Bloch's theorem of the form

$$\begin{pmatrix} u_n(\mathbf{x}) \\ v_n(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} u_{n,\mathbf{k}}(z) \\ v_{n,\mathbf{k}}(z) \end{pmatrix} e^{\mathrm{i}\mathbf{k}\cdot\mathbf{x}}, \tag{7}$$

where $u_{n,\mathbf{k}}(z)$ and $v_{n,\mathbf{k}}(z)$ are periodic functions with the same periodicity as the gap $\Delta_{\mathrm{LO}}(z)$ and are thus only periodic in the z-direction, which is reflected also in the single band index n.

We use the equilibrium state of the thermodynamic potential for the LO^1 phase as input to calculate $\alpha_\perp$ in order to determine the transition line between the LO^1 and LO^3 phase, see Fig.1(a).

Inhomogeneous Densities.— From solving the Bogoliubov-de Gennes equation in Eq.(6) we know the quasiparticle coherence factors u and v for the LO^1 phase. With these wavefunctions it is possible to



FIG. 3: Densities for the ^{6}Li atoms (a) and the ^{40}K atoms (b) scaled by the total density $n = n_\uparrow(0) + n_\downarrow(0)$ in the LO^1 phase as a function of position for the point denoted with a cross in Fig.1(a).

calculate the particle densities for the ^{6}Li and the ^{40}K atoms as a function of position. Namely, the densities are given by

$$n_\sigma(z) = \sum_{n,\mathbf{k}} \left\{ u^2_{n,\mathbf{k}}(z) N_\sigma(\hbar\omega_{n,\mathbf{k}}) + v^2_{n,\mathbf{k}}(z)[1 - N_{-\sigma}(\hbar\omega_{n,\mathbf{k}})] \right\} . \qquad (8)$$

We take the vector $\mathbf{k}$ to lie in the first Brioullin zone in momentum space, which is here infinitely large in the x and y direction and has size $2q$ in the z direction. In Fig.3 the inhomogeneous densities in the LO^1 phase are shown for the point denoted with a cross in the phase diagram Fig.1(a) and it can be seen that the density of the minority atoms follows the form of the Cooper-pair ansatz. In contrast, the density of the ^{40}K atoms has its maximum at the position where the Cooper pair has a minimum. In this way the system relaxes the frustration caused by the population imbalance, instead of phase separating, which occurs for a majority of ^{6}Li atoms [15, 16]. We find that the modulation in the density of the ^{6}Li atoms is about 10%, as can be seen in Fig.3(a), whereas the modulation for the majority atoms is much smaller. The density modulations can be visible in an experiment, for example using Bragg spectroscopy. A homogeneous superfluid does not show Bragg peaks and thus the appearance of any Bragg peak is a sign of inhomogeneities in the superfluid phase.

Discussion and Conlusion.— For the mixture of resonantly interacting ^{6}Li and ^{40}K atoms we completed the phase diagram, which contains inhomogeneous superfluid phases. These superfluid phases have a crystalline order, just as a solid, and are thus special kinds of supersolids. Usually a supersolid is defined as a phase with both long-range diagonal and off-diagonal order in the one-particle density matrix. The LO^1 and LO^3 only have diagonal order in the one-particle density matrix but there is both long-range diagonal and off-diagonal order in the two-particle density matrix. This means that the inhomogeneous superfluid is a supersolid phase, but not in the usual sense.

The results presented in this Letter were calculated making use of two approximations. The first one is the mean-field approximation, which is known to give good qualitative results. Although our results are only reasonable estimates quantitatively, we have great confidence that, as in the mass-balanced case, the phase diagram does not change qualitatively if screening and selfenergy effects are included. However, fluctuations can have a great effect on the density modulations. In order to observe a spatial periodicity in the densities true long-range order is most desirable. But due to fluctuation effects only algebraic long-range order will exist in the LO^1 phase [14, 31], making it more difficult to observe the periodicity.

The second approximation we make is that we take into account only a single magnitude of the lattice wavevector q. For the continuous transition from the normal gas to the superfluid state this is exact, whereas it is an approximation inside the superfluid phase. Deep in the superfluid phase the pairing field is expected to take a more complicated form and to depend on a range of wave vectors [13]. If in the phase diagram the chemical-potential difference, instead of the polarization, is on the x-axis all transitions take place close to the normal to superfluid transition. Therefore, it is not expected that taking into account a more complicated form for the gap will change the phase diagram qualitatively. Moreover, our calculation is variational and thus gives a conservative estimate of the supersolid region of the phase diagram.

In this Letter we presented results for a homogeneous system and we did not consider a confining trap. To take into account the trap for the atoms, a local-density approximation can be made. The trap felt by the atoms depends on the species of the atoms, which means that for two atom species two different traps have to be taken into account [19]. The trap can have an advantageous influence. Namely, if the periodicity of the Cooper pair wavefunction is pinned by the shape of the trap it becomes easier to observe density modulations.

This work is supported by the Stichting voor Fundamenteel Onderzoek der Materie (FOM), the Nederlandse Organisatie voor Wetenschaplijk Onderzoek (NWO).

[1] C.A. Regal *et al.*, Phys. Rev. Lett. **92**, 040403 (2004).
[2] M.W. Zwierlein *et al.*, Phys. Rev. Lett. **92**, 120403 (2004).
[3] M. W. Zwierlein *et al.*, Science **311**, 492 (2006).
[4] G. B. Partridge *et al.*, Science **311**, 503 (2006).
[5] Y. Shin *et al.*, Nature **451**, 689 (2008).
[6] K.B. Gubbels and H.T.C. Stoof, Phys. Rev. Lett. **100**, 140407 (2008).

[7] P. Fulde and R.A. Ferrell, Phys. Rev. **135**, A550 (1964).

[8] A.I. Larkin and Y.N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47**, 1136 (1964).

[9] Y. Liao, A.C. Rittner, T. Paprotta, W. Li, G.B. Partridge, R.G. Hulet, S.K. Baur, E.J. Mueller, Nature **467**, 567-569 (2010).

[10] C. Mora and R. Combescot, Phys. Rev. B **71**, 214504 (2005).

[11] N. Yoshida and S.-K. Yip, Phys. Rev. A **75**, 063601 (2007).

[12] L. Jiang and J. Ye, Phys. Rev. B **76**, 184104 (2007).

[13] A. Bulgac and M.M. Forbes, Phys. Rev. Lett. **101**, 215301 (2008).

[14] L. Radzihovsky, Phys. Rev. A **84**, 023611 (2011).

[15] K.B. Gubbels, J.E. Baarsma, and H.T.C. Stoof, Phys. Rev. Lett. **103**, 195301 (2009).

[16] J.E. Baarsma, K.B. Gubbels, and H.T.C. Stoof, Phys. Rev. A **82**, 013624 (2010).

[17] E. Wille *et al*, Phys. Rev. Lett. **100**, 053201 (2008).

[18] A.-C. Voigt *et al*, Phys. Rev. Lett. **102**, 020405 (2009).

[19] A. Trenkwalder, C. Kohstall, M. Zaccanti, D. Naik, A.I. Sidorov, F. Schreck, and R. Grimm, Phys. Rev. Lett. **106**, 115304 (2011).

[20] A. Ridinger, S. Chaudhuri, T. Salez, U. Eismann, D.R. Fernandes, K. Magalhães, D. Wilkowski, C. Salomon, and F. Chevy, EPJ D **65**, 223 (2011).

[21] C. Kohstall, M. Zaccanti, M. Jag, A. Trenkwalder, P. Massignan, G.M. Bruun, F. Schreck, and R. Grimm, Nature **485**, 615 (2012).

[22] R. Casalbuoni and G. Nardulli, Rev. Mod. Phys. **76**, 263 (2004).

[23] Y. Matsuda and H. Shimahara, J. Phys. Soc. Jpn. **76**, 051005 (2007).

[24] R. Casalbuoni, R. Gatto, N. Ippolito, G. Nardulli and M. Ruggieri, Phys. Lett. B **627**, 89 (2005)

[25] M. Mannarelli, K. Rajagopal and R. Sharma, Phys. Rev. D **73**, 114012 (2006).

[26] K. Rajagopal and R. Sharma, Phys. Rev. D **74**, 094019 (2006).

[27] N. D. Ippolito, G. Nardulli and M. Ruggieri, JHEP04, 036 (2007).

[28] M. Mannarelli, K. Rajagopal and R. Sharma, Phys. Rev. D **76**, 074026 (2007).

[29] J.M. Edge and N.R. Cooper, Phys. Rev. Lett. **103**, 065301 (2009).

[30] I. Zapata, F. Sols, and E. Demler, Phys. Rev. Lett. **109**, 155304 (2012).

[31] L. Radzihovsky and A. Vishwanath, Phys. Rev. Lett. **103**, 010404 (2009).